

# First Cenomanian record of insects in the Southern Hemisphere, with Perforissidae (Fulgoroidea) and Cupedidae (Coleoptera) from Southern Patagonia, Argentina



Julián F. Petrulevičius<sup>a, b, \*</sup>, Augusto N. Varela<sup>c</sup>, Ari Iglesias<sup>d</sup>, Alba B. Zamuner<sup>a, e, 1</sup>, Daniel G. Poiré<sup>c</sup>

<sup>a</sup> CONICET– Consejo Nacional de Investigaciones Científicas y Técnicas, Argentina

<sup>b</sup> División Palaeozoología, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, Paseo del Bosque s/n, La Plata 1900, Argentina

<sup>c</sup> Centro de Investigaciones Geológicas UNLP-CONICET, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, 1 N° 644, La Plata 1900, Argentina

<sup>d</sup> Instituto de Investigaciones en Biodiversidad y Ambiente INIBIOMA-CONICET, Universidad Nacional del COMAHUE, Quintral 1250, San Carlos de Bariloche 8400, Argentina

<sup>e</sup> División Paleobotánica, Facultad de Ciencias Naturales y Museo, Universidad Nacional de La Plata, Paseo del Bosque s/n, La Plata 1900, Argentina

## ARTICLE INFO

### Article history:

Received 2 August 2013

Accepted in revised form 24 May 2014

Available online

### Keywords:

Insecta  
Fulgoromorpha  
Archostemata  
Plant-insect interaction  
Mata Amarilla Formation  
Cenomanian  
Patagonia

## ABSTRACT

The discovery of the southernmost world record (49.5°S) of a Late Cretaceous insect fauna and plant-insect interactions is reported herein. The new locality is from the middle Cenomanian lacustrine deposits of the Mata Amarilla Formation belonging to the Austral foreland Basin, southwest Patagonia, Argentina. A first trip to the locality yielded few specimens of Coleoptera, among them an Archostemata: Cupedidae, and also a tiny specimen of Fulgoroidea: Perforissidae represented by *Aonikenkissus zamunerae* gen. et sp. nov. The Perforissidae were previously recorded from the Berriasian (?)–Barremian, Aptian, Albian, Turonian and Santonian in the Northern Hemisphere. The new species represents the first record of Perforissidae for South America and for the whole Southern Hemisphere. The new record suggests that the Perforissidae had a broader biogeographical distribution, and seem to have been cosmopolitan in the mid-Cretaceous. The Cupedidae reported herein is the first record for the Cenomanian and the first record of the genus *Zygadenia* in South America. Insect traces found in the same strata are made in an angiosperm palmatilobed leaf and correspond to a piercing and sucking type of damage. This type of damage is compared to recent stippling and to isolated piercings.

© 2014 Elsevier Ltd. All rights reserved.

## 1. Introduction

The discovery of the southernmost world record (49.5°S; palaeolatitude  $\approx 54^\circ$ ) of a Late Cretaceous insect fauna and plant-insect interactions is reported herein. The insects and traces were found in lacustrine deposits of the Mata Amarilla Formation that belongs to the Austral foreland Basin (=Magallanes Basin) in southwest Patagonia (Feruglio in Fossa Mancini et al., 1938; Leanza, 1972) (Fig. 1). This formation has a rich fauna of both marine invertebrates (Ihering, 1907; Wilckens, 1907; Feruglio, 1936; Riccardi,


1984a; 1984b; 2002; Riccardi and Rolleri, 1980; Riccardi et al., 1987; Griffin and Varela, 2012) and marine and continental vertebrates as crocodiles, turtles, theropods, plesiosaurs and sauropod dinosaurs (Goin et al., 2002; Lacovara et al., 2004; Novas et al., 2004a, 2004b, 2005, 2008; Cione et al., 2007; O’Gorman and Varela, 2010). It is also known for the richness in plants, especially unusual early angiosperm dominant floras (Berry, 1928, 1937; Frenguelli, 1953; Iglesias et al., 2007, 2009; Zamuner et al., 2004).

The Mata Amarilla Formation is considered as Cenomanian-Coniacian by Riccardi and Rolleri (1980) based on ammonites found in associated marine facies. Recently, precise U-Pb dating using laser ablation in zircons from continental facies from the middle section of the formation, gave 96.23  $\pm$  0.71 Ma in a tuff horizon about 28 meters above of the corresponding horizon where the insect and interactions were found (Varela, 2011; Varela et al., 2012a). On that basis, the new fossil record could be well

\* Corresponding author.

E-mail addresses: [levicius@fcnym.unlp.edu.ar](mailto:levicius@fcnym.unlp.edu.ar) (J.F. Petrulevičius), [varela@cig.museo.unlp.edu.ar](mailto:varela@cig.museo.unlp.edu.ar) (A.N. Varela), [ari\\_iglesias@yahoo.com.ar](mailto:ari_iglesias@yahoo.com.ar) (A. Iglesias), [poire@cig.museo.unlp.edu.ar](mailto:poire@cig.museo.unlp.edu.ar) (D.G. Poiré).

<sup>1</sup> Deceased in August 10, 2012.



**Fig. 1.** A, regional location map showing study area in dark gray. B, detailed map of study area. C, geological section of Mata Amarilla Formation (middle Cenomanian) at the studied locality. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)

restricted to the middle Cenomanian. The underlying Piedra Clavada Formation is well limited to the early/middle Albian by ammonoids (Piatnitzky, 1938; Riccardi et al., 1987) and palynomorphs (Archangelsky et al., 2008) with a transitional conformity between both units.

The entomofauna is composed by Fulgoroidea and Coleoptera. The specimen described herein belongs to Perforissidae Shcherbakov, 2007, a small and recently recognized extinct fulgroid family. The Perforissidae have 7 genera and were only known in the Cretaceous of the Northern Hemisphere (Shcherbakov, 2007; Peñalver and Szwed, 2010; Nel et al., 2012; Szwed et al., 2013). They are present in the Lower Cretaceous of Lebanese (Berriasian-Barremian; Szwed et al., 2013) and Burmese (Albian; Shcherbakov, 2007) ambers and Tsagan-Nur deposits (?Aptian; Shcherbakov, 2007), the Upper Cretaceous of New Jersey (Turonian; Shcherbakov, 2007), Taimyr (Santonian; Shcherbakov, 2007) and San Just (Albian; Peñalver and Szwed, 2010) ambers (see distribution in Fig. 1). Perforissidae is an enigmatic group concerning its evolutionary relationships within Fulgoroidea (Shcherbakov, 2007; Peñalver and Szwed, 2010). The Coleoptera are represented by fragmentary specimens. Among them it is remarkable one Cupedidae, that extends the distribution of this family in the Upper Cretaceous.

Insect traces were found in an angiosperm palmated leaf, corresponding to a piercing and sucking type of damage *sensu* Labandeira et al. (2007). Piercing and sucking fossil damage is discussed and compared to recent stippling. Stippling injuries are described from a recent species of Cicadellidae (Auchenorrhyncha) hosting Bignoniaceae leaves.

Mesozoic localities with insects become more and more recorded in last years in Argentina where several outcrops from Patagonia and Cuyo (west-central) geographic regions preserve some prospective insect faunas from the Middle–Upper Jurassic (Andrade de Moray and Genise 2005; Petrulevičius 2007) and the Cretaceous (Petrulevičius and Nel, 2003; Petrulevičius et al., 2005, 2010).

Upper Cretaceous insect localities are less common than those from the Lower Cretaceous in the world. In the same way the number of collected insects in the Upper Cretaceous is scarce in relation to the abundant material known from the Lower Cretaceous (Zherikhin, 2002). Even more, the Cenomanian localities with compressions of insects are few and only two were known outside Eurasia: the Canadian Knob Lake (Carrano, 2011) and the Lebanese Nammoura (Nel et al., 2004). In this sense, the discovery of a new locality in the middle Cenomanian of the South Hemisphere increases the interest for the understanding of the insect diversity and biogeography in the Upper Cretaceous.

## 2. Study area, material and methods

The insects and plant-insect interactions were recovered from the Cerro Waring Locality (Fig. 1, S 49° 31' and W 71° 30'00B4), from the Mata Amarilla Formation, in a fossiliferous level situated stratigraphically above the MAL and MEL plantiferous levels from Iglesias et al. (2007) and 28 m below the dated tuff level (Fig. 1). The specimens were found along with freshwater fish scales, fern, conifer and angiosperm remains, interpreted as a lacustrine palaeoenvironment along diverse vegetation in a continental lowland area (Varela, 2011). The insects and leaves with traces are kept in the collection of the Museo Provincial Padre Manuel Jesús Molina, Río Gallegos, Santa Cruz Province, Argentina with acronym MPM.

For wing venation nomenclature the system proposed by Kukalová-Peck (1983, 1991) and Dworakowska (1988) is followed, also used in previous works of the first author (JFP) dealing with Fulgoroidea (Petrulevičius, 2005; Solórzano Kraemer and

Petrulevičius, 2007). The forewing venation of Perforissidae is reduced. There are four main visible veins in the remigium interpreted as subcosta posterior plus radius anterior (ScP+RA), radius posterior (RP) media posterior (MP) and cubitus anterior (CuA). In some genera, and this is the case for the described species, the MP is fused with the CuA in the proximal part of the wing. More recent interpretations of wing venation of Fulgoroidea, like that of Nel et al. (2012), coincide in general with the Kukalová-Peck's system and with the present contribution specifically in the fusion of MP with CuA. They differ with Kukalová-Peck's system in the path of some veins like in the cubital system but the explicitly of this details exceeds the aims of this paper.

The Perforissidae and fossil leaf (Fig. 2A) were photographed with a Nikon D5000 camera and detail of the traces with an Olympus microscope BX51 with an Olympus epifluorescent unit and a Endow GFP BP filter (440–640 nm), connected to an Olympus Evolt E-330 camera, from the Instituto de Fisiología Vegetal (INFIVE) from La Plata National University (UNLP). The Perforissidae was drawn with a camera lucida attached to a Wild M5 stereomicroscope. The damaged recent leaves of *Tecoma stans* (Linnaeus) Juss. ex Kunth (Bignoniaceae) and the leafhoppers of the genus *Rhabdotalebra* Young (Cicadellidae: Typhlocybinae: Alebrini), producers of the stippling, were collected on May 15, 2012 at La Plata (Buenos Aires, Argentina; S 34°55' and W 57°57'). The insects correspond to an undescribed species (Catalano et al., in prep.) found to damage a new host plant for the genus *Rhabdotalebra* (Catalano et al., 2010). Photograph of the whole leaf was taken just after collection (Fig. 2B) with a camera Nikon D5000. In an attempt to better photograph the insect on the leaf the latter was frozen at –18 °C for about 5 minutes. See the freezing of the areolas that became black at the apex of the leaf (Fig. 2B detail). The damages became brown forming a ring sometimes open like a horseshoe (Fig. 2C details). After freezing, the leaf together with the insect were exposed to 4 °C to take the detailed photo of the Fig. 2B. Another leaf was dehydrated after putting it between card papers for a week (Fig. 2C) and photographed in details under a Leica microscope DM2500. The leaves and insects were collected on May, 14th (autumn) from La Plata, Buenos Aires, Argentina. Specimens are housed in the Entomological collection (Box No. 2012) of the Museo de La Plata (MLP), La Plata, Argentina.

## 3. Systematic palaeontology

Fulgoromorpha Evans 1946

Fulgoroidea Latreille 1807

Family Perforissidae Shcherbakov, 2007

Genus *Aonikenkissus* gen. nov. Petrulevičius, Varela, Iglesias and Poiré


*Derivation of name.* From the Aonikenk, ancient inhabitants of south Patagonia; and *Issus*, recent genus of the Issidae.

*Type species.* *Aonikenkissus zamunerae* gen. nov. et sp. nov., by monotypy and present designation.

*Diagnosis.* Tegmina gibbous with four main stems and distal branching (5–7 (?) apical cells); tegmen dark with pale-margined veins; costal area distally narrow; ScP+R, MP and CuA1+2 sinuous and subparallel in all their trajectory; crossveins r–m, m–cua and icua aligned; MP not stalked with ScP+R; CuA+MP stem as long as basal cell; CuA forked; basal part of MP near to CuA.


*Aonikenkissus zamunerae* sp. nov. Petrulevičius, Varela, Iglesias and Poiré

Figs. 3–4



**Fig. 2.** Insect damages in fossil and recent leaves. A, pentalobate leaf from Mata Amarilla Flora (MA106 angiosperm leaf morphotype; Iglesias et al., 2007), MPM-PB-1462. Note in the detail, the damages with a central undamaged area and surrounding the damaged ring, image with epifluorescence technique. B and C, recent leaves of *Tecoma stans* with stippling made by an undescribed species of *Rhabdotalebra* (Cicadellidae: Thyphlocybinae: Alebrini). B, fresh leaf, note in the detail the changes made to the damage after freezing the leaf, the producer was photographed in cold conditions (4 °C). After freezing, the habitus of the damage became similar to that of the fossil. C, dehydrated leaf, above, detail of a group of circular and horseshoe damages, right, detail of the individual injuries (marked by arrows) of a damage. Scale bars represent 3 mm in A, 0.2 mm in detail, 2 mm in B, 1 mm in detail, 4 mm in C, 0.2 mm in detail above, 0.5 mm in right detail.

**Derivation of name.** Named after Alba B. Zamuner, one of our co-authors, great colleague and outstanding paleobotanist from the Paleobotany Division at La Plata Museum. She passed away in August 10, 2012.



**Fig. 3.** *Aonikenkissus zamunerae* gen. et sp. nov., holotype MPM-PI-14932. Photograph of the tegmina. Scale bar represents 0.5 mm.

**Material.** Fragment of body and fore wing, MPM-PI-14932.

**Diagnosis.** As for the genus.

**Age and Outcrop.** Middle Cenomanian (96.23 ± 0.71 Ma), Mata Amarilla Formation, Cerro Waring, South Patagonia, Argentina.

**Description.** From fore wing; tegmina 3 mm long 1.2 mm wide; anterior areas not broadened; cells darkened with veins pale-margined (Fig. 3–4A); costal margin convex but slightly concave at the basal half (slightly sinuate), gibbous about mid-length; four main branches, i.e., ScP+RA, RP, MP, and CuA; MP+CuA stem about as long as basal cell, 0.23 mm long; arculus short; ScP+RA, RP and MP sinuous and running almost parallel; vein CuA not curved as three previous veins and running parallel to the claval fold; CuA bifurcating 0.57 mm basal to the crossveins, CuA1+2 slightly sinuous and fusing (?) with MP for a distance; CuA3+4 not changing direction from CuA; AA long running parallel to the claval fold; AP just preserved; posterior wing margin preserved only basal to the joining of AA with AP; RA with one branch;




Fig. 4. *Aonikenkissus zamunerae* gen. et sp. nov., holotype MPM-PI-14932. A, drawing of the tegmina. B, interpretation of wing venation. Scale bars represent 0.5 mm.

RP with one branch; MP with two (?) apical branches; CuA with three (or four ?) branches; 5–7 (?) apical cells; crossveins ra-rp, rp-m and mp-cua aligned; consequently to the alignment of these crossveins, the ScP+RA, the crossveins and the CuA1+2 result in a particular z-like drawing; crossvein cua1+2-cua3+4 (icua) distal to fusion of MP with CuA (? mp-cua); AP and AA+AP seems to be in straight line; crimped marginal membrane visible between RP to CuA.

**Discussion.** Main veins of the specimen are interpreted as ScP+RA, RP, and MP+CuA that separates into MP and CuA (Fig. 4B). Presence of a distal bifurcation of the fourth vein makes us to interpret it as the CuA basally fused to the MP. There is no stalk between the ScP+RA and RP; they are separated at the basal cell.

The specimen is included in Perforissidae because it shares all characters present in the fore wing of the family, i.e., wings small to minute (3–5 mm), tegmina narrow with four main stems (ScP+RA, RP, MP and CuA) and little distal branching (6–10), and a basal fusion of MP with CuA. The new genus shares some characters with *Perforissus* from the Late Cretaceous of New Jersey (Shcherbakov, 2007), i.e., tegmina gibbous about midlength; dark tegmen with pale-margined veins, a distally narrow costal area; MP+CuA stem equal than basal cell; ScP+RA, RP and MP sinuous and subparallel basal to crossveins; but differs with *Perforissus* in RP not stalked with ScP+RA; CuA forked and the crossveins ra-rp, rp-m, m-cua

aligned. In *Aonikenkissus* nov. gen. the veins themselves are not visible, but only the pale marginal area surrounding them. In *Perforissus*, as it is included in amber, it is possible to see the pale marginal area and the veins running, predominantly, at equal distance from both margins of it (see figs. 1, 3–5, 7 in Shcherbakov, 2007). Veins are interpreted in *Aonikenkissus* nov. gen. based on *Perforissus* model (Fig. 4B). The unique other genus with RP not stalked with the ScP+RA is *Cixitettix*. It differs from the new genus in having a single CuA, MP+CuA stem longer than basal cell and the tegmina widened to 2/3. Shcherbakov (2007) proposed two subfamilies within the Perforissidae. The definition of the taxa includes characters from the wing and hind leg tibia and tarsomere morphology (Shcherbakov, 2007). The wing character defining the Perforissinae, which is the costal margin gibbous at midlength (Shcherbakov, 2007; Peñalver and Szwed, 2010; Szwed et al. 2013) is also present in three genera of the other subfamily, the Cixitettiginae: *Foveopsis* (Shcherbakov, 2007), *Iberofoveopsis* (Peñalver and Szwed, 2010), and *Aafrita Szwed and Azar* (2013). In accord to this, the subfamilial categories have to be confirmed or redefined in the future for the Perforissidae. Within the family, the new specimen can be well placed in a new genus because a unique set of characters respect to the other genera, i.e., tegmina gibbous about midlength; tegmen dark with veins pale-margined; costal area distally narrow; ScP+RA not stalked with RP; CuA stem equal

**Table 1**  
Comparison of *Aonikenkissus* gen. nov. with the other genera of the family. Selected characters.

Characters \ genera	<i>Tsaganema</i>	<i>Cixitettix</i>	<i>Foveopsis</i>	<i>Iberofoveopsis</i>	<i>Aafrita</i>	<i>Aonikenkissus</i> gen. nov.	<i>Perforissus</i>	<i>Cretargus</i>
Teeth in hind leg pectens	Metatibia 1st tarsus 2nd tarsus	Uniserial				¿?	Triserial Biserial no	Biserial Uniserial
Tegmen color pattern		Contrasting pale and dark pattern	Unicolor dark			Cells darkened, veins pale margined		Pale with dark markings
Tegmen shape		Parallel sided	Widened to 2/3	Gibbous at 1/2				
Length MP+CuA stem versus basal cell		Equal	Longuer	Equal	Shorter		Shorter	Shorter
ScP+RA +RP stalk		Short	No	Long	Short		Long	
CuA		Normally unforked	Unforked	Forked		No	Unforked	
Wing length versus crossveins to apex length		~1/3		~1/4		~1/5		¿?

than basal cell; CuA forked; basal part of MP near to CuA. *Aonikenkissus* nov. gen. has an unique character within the family: the forewing has the crossveins ra-rp, rp-m, and the m-cua aligned. Differences and similarities with the other genera of Perforissidae are detailed in Table 1.

Coleoptera Linné, 1758  
Archostemata Kolbe, 1968  
Cupedidae Laporte, 1836  
Notocupedini Ponomarenko, 1968  
Genus *Zygadenia* Handlirsch, 1906

*Zygadenia* sp. 1  
Fig. 5A

**Material.** MPM-PI-14716.

**Age and Outcrop.** Middle Cenomanian, Cerro Waring, South Patagonia, Argentina.

**Remarks.** The elytrum is fragmentarily preserved but shows the typical venation and ornamentation of the family Cupedidae (see Ponomarenko, 1966, 2000, 2006). The specimen MPM-PI-14716 (Fig. 5A) could be attributed to the genus *Zygadenia* Handlirsch, 1906 (see Ponomarenko, 1966, 2000) because it has a distinct venation with two rows of cells between main veins, the main veins of the elytrum sharply distinct from the intercalaries forming large quadrangular cells, and the main veins seem to coalesce before the apex. Handlirsch (1906) proposed the generic name *Zygadenia* for the isolated elytra of Cupedidae with certain characters. Ponomarenko (2000) synonymised *Notocupes* Ponomarenko, 1968 (Ponomarenko, 1968) with *Zygadenia*, but it could be a better practice to use the name *Zygadenia* for isolated elytra (Ponomarenko and Ren, 2010). The specimen seems to be a new species but it is not described for the moment because of its fragmentary condition and the possibility of finding new material in the future.

Coleoptera indet.  
Fig. 5B–J.

**Material.** MPM-PI-14717, MPM-PI-14718, MPM-PI-14719, MPM-PI-14720, MPM-PI-14721, MPM-PI-14722, MPM-PI-14723, MPM-PI-14724, MPM-PI-14725.

**Remarks.** From the same strata were recorded several disarticulated insects attributed to Coleoptera, most of them correspond to elytra but also other parts as abdominal ventrites. Isolated elytra from the Cretaceous usually could not be assigned to familial level (Ponomarenko, 2006). Only the Archostemata Cupedidae Laporte, 1836 could be attributed without doubts and sometimes to generic level (Ponomarenko, 2006). There are 8 different morphotypes of elytra that could not be assigned certainly to any family of Coleoptera. One specimen has longitudinal punctate rows

(Fig. 5B), others have uniformly punctate grooves (Fig. 5C–D) and the dominant morphology has neither veins nor cells (Fig. 5E–I). There is also an isolated abdomen of Coleoptera MPM-PI-14725 (Fig. 5J) that has five ventrites with the metacoxae cavities not dividing the ventrite 1 and an apparently rugose texture; characters that coincide with archostematan morphology, but the specimen is too incomplete to be assigned to any family.


#### 4. Palaeobiogeography

*Aonikenkissus zamunerae* gen. nov. sp. nov. significantly enlarges the geographic distribution of Perforissidae (Fig. 6) being the southernmost species and the first record in the Southern Hemisphere. The Perforissidae are not represented in the Northern Hemisphere during the Cenomanian; but there are Upper Cretaceous records in the Santonian of Taimyr and the Turonian of North America (Fig. 6). In this context the family distribution could be inferred as widespread at least in the mid-Cretaceous; also considering that the taxon is under-sampled because of its small size and the few paleoentomologists in the Southern Hemisphere. The latest hypothesis could explain also the first record of the archostematan genus *Zygaenia*, very well represented in the Jurassic-Lower Cretaceous of Eurasia and incipiently recorded in the South Hemisphere as in the Jurassic of Australia (Martin, 2010) and the present record. The gap in the record of Cupedidae in the Cenomanian (Ponomarenko, 2002; Soriano and Delclòs, 2006), is filled by the present record and possibly due to the scarcity of Upper Cretaceous localities with beetles, mainly three in the world (Ponomarenko, 2002).

In the Figure 6 are plotted all the records of Perforissidae in the Cretaceous and the Upper Cretaceous record of Cupedidae. In the same Figure it is seen that the record of Perforissidae is clearly related to amber localities, whereas the record of Cupedidae is related to compression localities. Among all the localities only two, the New Jersey amber and Mata Amarilla have both taxa, Perforissidae and Cupedidae (circles and squares in Fig. 6). The surprising first insects recorded in Mata Amarilla and the fact that the locality bears one of the southernmost entomofauna in the world for the Late Cretaceous, show that the sequences at the Austral Basin in Argentinean Patagonia are very interesting and promising.


#### 5. Perforissidae paleoecology and plant-insect interactions

In the case of Perforissidae it is difficult to make actualistic ecological inferences, not only because it is an extinct family, but also because it has obscure phylogenetic relationships within the Auchenorrhyncha and more precisely the Fulgoroidea. But due to the biogeographic pattern, rarity of the taxon and age, it results very interesting to discuss about the possible host plants and climatic conditions inhabited by this family. It is possible to



**Fig. 5.** A, *Zygadenia* sp. 1, MPM-PI-14716. B– I, Coleoptera indet., elytra. B, MPM-PI-14717. C, MPM-PI-14718. D, MPM-PI-14719. E, MPM-PI-14720. F, MPM-PI-14721. G, MPM-PI-14722. H, MPM-PI-14723. I, MPM-PI-14724. J, Coleoptera indet., ventral view of abdomen, MPM-PI-14725. Scale bars represent 2 mm in A and I, 1 mm in B–H and J.

CRETACEOUS	Upper	Maastrichtian	70.6	
		Campanian	83.5	
		Santonian	85.8	<i>Cretargus</i> <i>Cixitettix</i>
		Coniacian	88.6	
		Turonian	93.6	<i>Perforissus</i> Cupedidae
		Cenomanian	99.6	<i>Aonikenkissus</i> gen. nov. Cupedidae
		Albian	112	<i>Foveopsis</i> <i>Iberofoveopsis</i>
		Aptian	125	<i>Tsaganema</i>
	Lower	Barremian	130	
		Hauterivian	133.9	<i>Aafrita</i>
		Valangian	140.2	?
		Berriasian	145.5	



**Fig. 6.** World map with the localities and geological periods recording the Perforissidae (circles) and the Upper Cretaceous Cupedidae (squares). Geological periods with Perforissidae record in gray. Amber localities in color, Compressional ones in white. 1, Taimyr amber; 2, New Jersey amber; 3, Kyzyl-Syr; 4, Cerro Waring (Mata Amarilla Formation); 5, Burmese amber; 6, San Just amber; 7, Tsagan-Nur; 8, Lebanese amber. (For interpretation of the references to colour in this figure legend, the reader is referred to the web version of this article.)



indirectly restrict and hypothesize the life style and interactions of the group in relation to the physical environment and floral aspects of the taphocenosis of the concerned strata, and from some morphological features of the insect that permit some discussions.

Mata Amarilla Formation is known for their well preserved compression floras and petrified forests. These Cretaceous forests were dominated by Podocarpaceae (Zamuner et al., 2004; Iglesias et al., 2007), associated to both coastal–littoral marine and lowland environments (Varela, 2011). Understory and remaining flora, as the flora associated to the Perforissidae fossiliferous strata (Fig. 1), were mostly herbaceous or low shrubs dominated by ferns and primitive angiosperms (Iglesias et al., 2007, 2011). The angiosperm leaves associated to the perforissid have a relative high morphological diversity with broad and entire margins related to warm and wet environments; although brachyphyllous conifers are also present (Iglesias et al., 2007). The record of broad leaves with damages adds the possibility to maintain leafhopper-like animals related to them.

The sensory pits retained in the adults of some Fulgoroidea are presumed hygroreceptors (Shcherbakov, 2007). Although there is no empirical demonstration about the function of these organs (Bräunig et al., 2012), they are usually related to extreme xerophiles or hygrophiles, and they are also found in presence of strong seasonal variation (Shcherbakov, 2007). In the same way, Gnezdilov (2012) pointed that species with sternal sensory pits in the imago inhabit arid biotopes. So, the multiple presence of pits in Perforissidae could indicate that they inhabited biotopes with highly variable or extreme humidity, possibly coastal–littoral environments (Shcherbakov, 2007). After the geological reconstruction (Varela et al., 2011), the inhabited environment of *Aonikenkissus zamunerae* in Mata Amarilla Formation corresponds to a lowland area, probably a coastal lake; and in agreement to the regional reconstruction is relatively far (more than 200 km) from the sea (Varela, 2011; Varela et al., 2013).

These animals depend on plants to feed, but up to date, there are not evidence about their diet preferences. Perforissids have in general a morphology of the rostrum comparable to that of recent plant/leafhoppers and suitable to produce similar plant damages like piercing and sucking (Fig. 2). Based on the whole body and 3D preservation in amber, Shcherbakov (2007) interpreted the Perforissidae as an early attempt to construct leafhopper-like forms from a planthopper group. This “cicadelization” (attempt to construct cicadellid forms; Shcherbakov, 2007) could imply a leafy diet for these insects as for the recent Cicadellidae. Damages by recent leafhoppers can be recognized because they introduce their stylets in the mesophyllous of the leaves and suck their fluids producing different foliage injuries, like stipples, burning (Backus et al., 2005) and isolated piercings (Backus et al., 1988). All Auchenorrhyncha have intracellular penetration style contrary to the Sternorrhyncha that have intercellular penetration (Backus et al., 2005; Tjallingii, 2006). Different species use different feeding strategies as cell rupturing and salivary sheaths, the latter one used by most of aucherrhynchans (Backus et al., 2005). These two strategies could be differentiated from the surface of Recent leaves. Cell rupturing in the case of stippling produces different white “drawings” as stars, horseshoes or rings (See Fig. 2B–C; also Cranshaw, 2010; Brentassi et al., 2010). Salivary flanges (and underlying sheaths) could be visible but using a previous staining method (Backus et al., 1988).

Angiosperm leaves recorded at Mata Amarilla Formation show leaf damages that can be unequivocally assigned to diverse arthropod actions. The Fig. 2A shows possible piercing and sucking damages in a palmatilobed angiosperm leaf, similar to DT46 leaf damage type from Labandeira et al. (2007).

Several recent species of Thyphlocybinae produce stipples resulting in different white “drawings” in angiosperm leaves. In Fig. 2B and 2C are shown the damages produced by an undescribed species of the genus *Rhabdotalebra* (Cicadellidae: Thyphlocybinae: Alebrini; Catalano et al., in prep.) formed by rings or horseshoes. These damages are grouped around the principal veins, producing stippling sometimes in high concentration in *Tecoma stans* (Bignoniaceae) leaves. Each ring is the result of several injuries (Fig. 2C details). Individual injuries forming horseshoe to ring forms are in turn aggregated in patches, or linear or zigzag patterns (Petrulevičius, pers. obs.). So, the damages by this species have inclusive spatial levels of complexity.

When freezing the leaf of Fig. 2B, the white stipples turned to brown. The central part surrounded by the ring remained green, as well as the surrounding surface (not injured) of the leaf (Fig. 2C details). The result unexpectedly and interestingly is quite similar to the fossil damages found in the Cretaceous leaf (Fig. 2A). Considering this result it was interpreted that in some damages like the DT46 of Labandeira et al. (2007) and that on the leaf here described stipples have not to be discarded as a possible explanation. Then it is considered the possibility of production of fossil piercing and sucking damages (sensu Labandeira et al., 2007) since the middle Cretaceous in angiosperm leaves, either by one central piercing (and sucking by a salivary sheath) or by several injuries ring like performed. Coincidentally, stipples are known to be produced in recent times by the thyphlocybines which have a pantropical distribution and seem to have arisen during the Cretaceous (Dietrich, 1999). Obviously, it is not possible to know what kind of damage produces our insect, the plant organ or even the kind of plant on which it feeds, brachyphyllous or low herbaceous, as interpreted by Shcherbakov (2007) or angiosperm understory forest and low shrubs with broad leaves which are the dominant plants in Mata Amarilla (Iglesias et al., 2007). The record of damages is restricted for the moment in Mata Amarilla to an angiosperm broad leaf possibly belonging to a forest understory plant (Fig. 2A).

Shcherbakov (2007) referred as possible causes of Perforissidae extinction the competition with the Cicadellidae or the fact that they were trophically associated with earliest angiosperms or “proangiosperms” (Shcherbakov, 2007; term referring in part ? to gymnosperm Hirmerellaceae), lineages not surviving (for the authors) into the Cenozoic (Shcherbakov, 2007; Peñalver and Szewdo, 2010). Although almost all angiosperms known in the Mata Amarilla Fm. are extinct by the latest Cretaceous, it has to be taken into account that Perforissidae seem to be strongly underrepresented in the fossil record as compressions by taphonomical and collection biases (due to its small size), so its extinction still could be somewhere before or after the Cretaceous/Paleogene (K/Pg). Besides, one of the groups considered extinct in the Cretaceous, the Hirmerellaceae producing *Classopollis* pollen surpassed the Upper Cretaceous, and are found in the Paleocene of Patagonia (Archangelsky and Romero, 1974; Pöthe de Baldi, 1984; Quattrocchio and Ruiz, 1999) and Northwest Argentina (Petrulevičius, 1996; Quattrocchio and Del Papa, 2000). In this sense the K/Pg boundary seems not to be critical for these plants nor for other insect groups as the Odonata that recognize a crisis through changes of the morphology/taxonomy between the Early and Late Cretaceous and not between the K/Pg boundary (Nel et al., 2010). Studies on insects and plant-insect interactions are needed for the Southern Hemisphere to better compare how stronger were environmental changes between the Early and the Late Cretaceous and K/Pg boundary transition.

## 6. Discussion about climate

The Mata Amarilla Formation represents a continental low-land area eventually flooded by epeiric seas, although only the lower and the upper sections of Mata Amarilla Formation were developed in a mixed littoral environment (Varela et al., 2011). Multiple stacked paleosoils are recognized throughout the Mata Amarilla Formation. Although the middle section preserves the most characteristic paleosoils, where the insect remains were found, associated to margins of shallow lakes (Fig. 1). The histosols, vertisols, alfisols and inceptisols recognized at the Middle Section of the unit indicate temperate warm climate, i.e., mean annual precipitations ~1000 mm, and mean annual temperatures close to 20 °C (Varela et al., 2012b). Geological indicators in the soils, detected marked seasonal dry periods, explained by not evenly distributed rains through the seasons (Varela et al., 2012b).

The angiosperm leaf characters as margin type and size described for the Mata Amarilla Formation flora, also are related to warm and humid climates (Iglesias et al., 2007). On the other hand, the presence of defined wood rings in most of the conifer trunks preserved in the Lower Section of the unit, allows inferring growth rate seasonality linked to some kind of seasonality in the water supply (Zamuner et al., 2004).

Cretaceous Cupedidae could be correlated mainly with warm climates (Soriano and Delclòs, 2006) but some records are outside to the subtropical climate area proposed by Spicer et al. (1994). This is the case of the Siberian Kyzyl-Syr locality, unique previous record of the genus *Zygadenia* (Ponomarenko, 1969) in the Late Cretaceous (Soriano and Delclòs, 2006). The other record of the family in the Late Cretaceous is from the Turonian New Jersey amber (Fig. 6). The record of Cupedidae in the Late Cretaceous is rare, in contrast to the Early Cretaceous (4 vs. 57 species, Soriano and Delclòs, 2006 and this work).

The climate inferred by geological indicators is indicative of biotopes with highly or extremely variable humidity, the same seems to be marked by grow rings in the fossil woods. This climate is probably related with the presence of the multiple sensory pits of the Perforissidae.

## 7. Concluding remarks

Perforissids are rare in the fossil record and mainly recorded in amber, excepting only two occurrences in compressional lacustrine deposits: the Mongolian record (Ponomarenko and Ren, 2010) and the specimen described here. This fact could be related to the tiny size of the specimens that are more suitable to be trapped in resins and not normally preserved nor recognized in compressions. The Argentinean specimen represents the first Perforissidae recorded for the whole Southern Hemisphere. This discovery confirms that the family was widely distributed in both hemispheres during the mid-Cretaceous.

Coleoptera is the most abundant group of the new insect fauna. The discovery of a Cupedidae fills the Upper Cretaceous world gap for this family in the Cenomanian and gives the first record of the genus *Zygadenia* in South America.

Insect traces found in Mata Amarilla are recorded in an angiosperm broad leaf and correspond to a piercing and sucking type of damage. This type of damage was compared, via experimentation, with recent stipples, arriving to the possibility of production either by one central piercing (and sucking by a salivary sheath) or by several ring-like injuries.

## Acknowledgments

This research was supported by grants from the National Research Council of Argentina (CONICET): PIP 1016, PIP 6393, and PIP 0377. We are indebted to the Agencia Nacional de Promoción Científica y Tecnológica de Argentina (ANPCyT) by grant PICT-2012-1555 and to the National Science Foundation of USA (NSF) by grant DEB-0919071 for technical support. We thank the Culture Secretary of Santa Cruz province for permits. DGP, ABZ, ANV and AI thank to: S. Richiano and C. Koefoed, for exceptional field assistance; to the landlords Waring, Rolón, Bertotti, and Pichinini for their friendship and field assistance; to the “Comisaría, Escuela and Comisión de Fomento” from Tres Lagos, and M. Ferrari for logistics at the field work. We also thank J. Szewo and A. Ponomarenko for valuable discussions about Fulgoroidea and Coleoptera, respectively; and the two anonymous reviewers that strongly enhanced this contribution.

## References

- Archangelsky, S., Romero, E.J., 1974. Polen de gimnospermas (Coníferas) del Cretácico Superior y Paleoceno de Patagonia. *Ameghiniana* 11, 217–236.
- Archangelsky, A., Archangelsky, S., Poiré, D.G., Canessa, N.D., 2008. Registros palinológicos en la Formación Piedra Clavada (Albiano) en su área tipo, provincia de Santa Cruz, Argentina. *Revista del Museo Argentino de Ciencias Naturales* 10, 185–198.
- Andrade de Moray, M., Genise, J.F., 2005. An association of fossil larval tubes and head capsules of Chironomidae (Diptera) from the Jurassic (Callovian-Oxfordian) Cañadon Asfalto Formation, Patagonia, Argentina. Abstracts volume. III International congress of palaeontomology, II international meeting on palaeoarthropodology, and II world congress on amber and its inclusions (Fossils X3), Pretoria, p. 9.
- Backus, E.A., Hunter, W.B., Arne, C.N., 1988. Technique for Staining Leafhopper (Homoptera: Cicadellidae) Salivary Sheaths and Eggs Within Unsectioned Plant Tissue. *Journal of Economic Entomology* 81, 1819–1823.
- Backus, E.A., Serrano, M.S., Ranger, C.M., 2005. Mechanisms of hopperburn: An Overview of Insect Taxonomy, Behavior, and Physiology. *Annual Review of Entomology* 50, 125–151.
- Berry, E.W., 1928. Tertiary fossil plants from the Argentine Republic. *Proceedings of the United States National Museum* 73 (22), 1–27.
- Berry, E.W., 1937. An Upper Cretaceous flora from Patagonia. *Johns Hopkins University – Studies in Geology* 12, pp. 11–31.
- Bräunig, P., Krumpal, K., Baumgartner, W., 2012. Sensory pits - Enigmatic sense organs of the nymphs of the planthopper *Issus coleoptratus* (Auchenorrhyncha, Fulgoromorpha). *Arthropod Structure & Development* 41, 443–458.
- Brentassi, M.E., Catalano, M.I., Paradell, S., Marino de Remes Lenicov, A.M., 2010. Caracterización de *Typhlocybella maidica* (Hemiptera: Cicadellidae) y descripción del daño producido en plantas de maíz y gramíneas asociadas en la Argentina. *Revista de la Sociedad Entomológica Argentina* 69, 57–64.
- Carrano, M., 2011. PaleoDB collection 119438, Redmond No. 1, Knob Lake District (Cretaceous of Canada). In: Karr, J., Clapham, M. (Eds.), *Paleobiology Database*. [http://paleodb.org/cgi-bin/bridge.pl?action=basicCollectionSearch&collection\\_no=119438](http://paleodb.org/cgi-bin/bridge.pl?action=basicCollectionSearch&collection_no=119438).
- Catalano, M.I., Paradell, S.L., de Remes Lenicov, A.M.M., 2010. Revision of the genus *Rhabdotalebra* Young (Hemiptera: Cicadellidae: Typhlocybinae: Alebrini) and description of two new species from Argentina. *Zootaxa* 2601, 53–60.
- Catalano, M.I., Paradell, S.L., Fraga, S.V., Petrulevičius, J.F., In prep. A new species the genus *Rhabdotalebra* from Argentina (Hemiptera: Cicadellidae: Typhlocybinae).
- Cione, A.L., Gourie, S., Goin, F., Poiré, D., 2007. *Atlantoceratodus*, a new genus of lungfish from upper Cretaceous of South America and Africa. *Revista del Museo de La Plata. Paleontología* 10 (62), 1–12.
- Cranshaw, W., 2010. Colorado State University, Bugwood.org. <http://www.forestryimages.org/browse/detail.cfm?imgnum=5369839>
- Dietrich, C.H., 1999. The role of grasslands in the diversification of leafhoppers (Homoptera: Cicadellidae): a phylogenetic perspective. In: Warwick, C. (Ed.), *Proceedings of the Fifteenth North American Prairie Conference*. Natural Areas Association, Bend, pp. 44–48.
- Dworakowska, I., 1988. Main veins of the wings of Auchenorrhyncha. *Entomologische Abhandlungen, Staatliches Museum für Tierkunde Dresden* 52 (3), 63–108.
- Feruglio, E., 1936. Nota preliminar sobre algunas nuevas especies de moluscos del Supracretáceo y Terciario de la Patagonia. *Notas del Museo de La Plata. Paleontología* 1, 277–300.
- Feruglio, E., 1938. In: Fossa Mancini, E., Feruglio, E., Yussen De Campana, J.C. (Eds.), *Una reunión de geólogos de YPF y el problema de la terminología estratigráfica*, Boletín de Informaciones Petroleras 171, pp. 31–95.
- Frenguelli, J., 1953. La flora fósil de la región del alto Río Chalia en Santa Cruz (Patagonia). *Notas Museo de La Plata (paleontología)* 98, 239–257.

- Gnezdilov, V.M., 2012. A New Species of the Genus *Rhinogaster* Fennah (Homoptera, Fulgoroidea, Caliscelidae) from Southern Africa. *Entomological Review* 92, 93–96.
- Goin, F.J., Poiré, D.G., De La Fuente, M.S., Cione, A.L., Novas, F.E., Bellosi, E.S., Ambrosio, A., Ferrer, O., Canessa, N.D., Carloni, A., Ferigolo, J., Ribeiro, A.M., Sales Viana, M.S., Pascual, R., Reguero, M., Vucetich, M.G., Marensi, S., De Lima Filho, M., Agostinho, S., 2002. Paleontología y Geología de los sedimentos del Cretácico Superior aflorantes al sur del Río Shehuen (Mata Amarilla, Prov. de Santa Cruz, Argentina). In: *Actas. XV Congreso Geológico Argentino*, pp. 603–608.
- Griffin, M., Varela, A.N., 2012. Mollusk fauna from the Mata Amarilla Formation (lower Upper Cretaceous), southern Patagonia Argentina. *Cretaceous Research* 37, 164–176.
- Handlirsch, A., 1906–1908. Die fossilen Insekten und die Phylogenie der rezenten Formen. V.W. publ., Leipzig. Engelman, 1430 pp.
- Iglesias, A., Zamuner, A., Poiré, D.G., Lariestra, F., 2007. Diversity taphonomy palaeoecology of an angiosperm flora from the Cretaceous (Cenomanian–Coniacian) in Southern Patagonia, Argentina. *Palaeontology* 50, 445–466.
- Iglesias, A., Zamuner, A., Poiré, D.G., Varela, A., Richiano, S., Koefed, C., 2009. Albian–Campanian continuous record of compression floras in Tres Lagos, Austral Basin, Patagonia, Argentina. Reunión de Comunicaciones Anuales de la Asociación Paleontológica Argentina, Ciudad de Bs. As, pp. 50–51.
- Iglesias, A., Artabe, A.E., Morel, E.M., 2011. The evolution of Patagonian climate and vegetation, from the Mesozoic to the present. *Biological Journal of Linnean Society Special Paper* 103, 409–422.
- Ihering, H.Von, 1907. Les mollusques fossiles du Tertiaire et du Crétacé supérieure de la Argentine. *Anales Museo Nacional de Buenos Aires* 7, 1–611.
- Kukalová-Peck, J., 1983. Origin of the insect wing and wing articulation from the arthropodan leg. *Canadian Journal of Zoology* 61, 1618–1669.
- Kukalová-Peck, J., 1991. Fossil history and the evolution of hexapod structures. In: Naumann, I.D., Crane, P.B., Lawrence, J.F., Neilsen, E.S., Spradbery, J.P., Taylor, R.W., Whitten, M.J., Littlejohn, M.J. (Eds.), *The Insects of Australia. A textbook for students and research workers* 1, Second edition. Melbourne University Press, pp. 141–179.
- Labandeira, C.C., Wilf, P., Johnson, K.R., Marsh, F., 2007. Guide to insect (and other) damage types on compressed plant fossils (Version 3.0-Spring 2007). Smithsonian Institution.
- Laporte, F.L. de Castelnaud, 1836. Etudes Entomologiques, ou descriptions d'insectes nouveaux et observations sur la synonymie. *Revue Entomologique* 4, 5–60.
- Lacovara, K., Harris, J., Lammana, M., Novas, F., Martinez, R., Ambrosio, A., 2004. An enormous sauropod from the Maastrichtian Pari Aike Formation of southernmost Patagonia. *Journal of Vertebrate Paleontology* 24, 81A.
- Leanza, A.F., 1972. Andes Patagónicos Australes. In: Leanza, A.F. (Ed.), *Geología Regional Argentina*. Academia Nacional de Ciencias, pp. 689–706. Córdoba, 869 pp.
- Martin, S.K., 2010. Early Jurassic coleopterans from the Mintaja insect locality, Western Australia. *Acta Geologica Sinica, English Edition* 84 (4), 925–953.
- Nel, A., Azar, D., Martínez-Delclós, X., Makhoul, E., 2004. A new Upper Cretaceous species of *Chresmoda* from Lebanon a latest representative of Chresmodidae (Insecta: Polyneoptera inc. sed.): first record of homeotic mutations in the fossil record of insects. *European Journal of Entomology* 101 (1), 145–151.
- Nel, A., Nel, P., Petrulevičius, J.F., Prokop, P., Perrichot, V., Azar, D., 2010. The Wagner parsimony using morphological characters: a new method for palaeoecological studies. In: Nel, A., Petrulevičius, J.F., Azar, D. (Eds.), *Fossil insects, Special issue Annales de la Société Entomologique de France* (n. s.), 46 (1–2), pp. 276–292.
- Nel, A., Prokop, J., Nel, P., Grandcolas, P., Huang, Di-ying, Roques, P., Guilbert, E., Dostál, O., Szweido, J., 2012. Traits and evolution of wing venation pattern in paraneopteran insects. *Journal of Morphology* 273, 480–506.
- Novas, F.E., Cambiaso, A.V., Ambrosio, A., 2004a. A new basal iguanodontian (Dinosauria, Ornithischia) from the Upper Cretaceous of Patagonia. *Ameghiniana* 41, 75–82.
- Novas, F.E., Lecuona, A., Calvo, J., Porfiri, J., 2004b. Un terópodo del Cretácico Superior de la provincia de Santa Cruz. *Ameghiniana* 41, 58.
- Novas, F.E., Salgado, L., Calvo, J.O., Agnolín, F.L., 2005. Giant titanosaur (Dinosauria, Sauropoda) from the Late Cretaceous of Patagonia. *Revista del Museo Argentino de Ciencias Naturales, nueva serie* 7, 37–41.
- Novas, F.E., Ezcurra, M.A., Leucona, A., 2008. *Orkoraptor burkei* nov. gen. et sp., a large theropod from the Maastrichtian Pari Aike formation, Southern Patagonia, Argentina. *Cretaceous Research* 29, 468–480.
- O'Gorman, J.P., Varela, A.N., 2010. The oldest lower Upper Cretaceous plesiosaurs (Reptilia, Sauropterygia) from the southern Patagonia, Argentina. *Ameghiniana* 47, 447–459.
- Peñalver, E., Szweido, J., 2010. Perforissidae (Hemiptera: Fulgoroidea) from the Lower Cretaceous San Just amber (Eastern Spain). *Alavesia* 3, 97–103.
- Petrulevičius, J.F., 1996. Nueva localidad fosilífera para el Paleoceno Superior (Formación Maíz Gordo) del Noroeste argentino. *Resúmenes Congreso Paleógeno de América del Sur*, p. 20.
- Petrulevičius, J.F., 2005. First fossil record of Nogodinidae in South America; a planthopper from the upper Paleocene of Argentina. *Systematics and taphonomy. Palaeontology* 48, 299–308.
- Petrulevičius, J.F., 2007. A new species of Bittacidae sensu lato (Mecoptera) from the Callovian–Oxfordian: new Jurassic locality of insect body fossils from Patagonia, Argentina. Abstracts IV International Congress of Paleontology, II International Meeting on Palearthropodology, and III Amber International Congress (“Fossil X3”), Vitoria-Gasteiz.
- Petrulevičius, J.F., Nel, A., 2003. Oldest Petalurid dragonfly (Insecta: Odonata): A Lower Cretaceous specimen from South Patagonia, Argentina. *Cretaceous Research* 24, 31–34.
- Petrulevičius, J.F., Sallenave, S., Nel, A., 2005. Ambush predator guild of Backswimmers in the Cretaceous of Argentina. Abstracts volume III International Congress on Palaeoentomology, II International Meeting on Palearthropodology and III International Congress on amber (Fossils X3), Pretoria, p. 29.
- Petrulevičius, J.F., Nel, A., Sallenave, S., 2010. Notonectidae (Insecta, Heteroptera) from the Cretaceous of San Luis, Argentina. In: Nel, A., Petrulevičius, J.F., Azar, D. (Eds.), *Fossil insects. Special issue Annales de la Société Entomologique de France* (n. s.) 46 (1–2), Paris, pp. 271–275.
- Piatnitzky, A., 1938. Observaciones geológicas en el Oeste de Santa Cruz (Patagonia). *Boletín de Informaciones Petroleras* 165, 45–85.
- Ponomarenko, A.G., 1966. New beetles of the family Cupedidae (Coleoptera) from Mesozoic deposits of Transbaikalia. *Entomologicheskoe Obozrenie* 45, 138–143 (in Russian, translated in English in *Entomological Review* 45, 75–79).
- Ponomarenko, A.G., 1968. Archostematan beetles from the Jurassic of Karatau (Coleoptera, Archostemata). In: Rohdendorf, B.B. (Ed.), *Yurskie nasekomye karatau [Jurassic insects of Karatau]*. Akademiya nauk SSSR Otdelenie Obshchej Biologii, Moscow, pp. 118–138.
- Ponomarenko, A.G., 1969. Istoricheskoe razvitiye zhestkokrylykh–arkhostemat [Historical development of the Coleoptera - Archostemata]. *Trudy Paleontologicheskogo Instituta Akademii Nauk SSSR* 125, 1–240 [In Russian].
- Ponomarenko, A.G., 2000. Beetles of the family Cupedidae from the Lower Cretaceous locality of Semen, Transbaikalia. *Paleontological Journal* 34 (Supplement 3), S.317–S.322.
- Ponomarenko, A.G., 2002. 2.2.1.3.2. Superorder Scarabaeidea Laicharting, 1781. Order Coleoptera Linné 1758. THE BEETLES. In: Rasnitsyn, A.P., Quicke, L.J. (Eds.), *History of Insects*. Kluwer Academic Publishers, pp. 164–176.
- Ponomarenko, A.G., 2006. On the types of Mesozoic archostematan beetles (Insecta, Coleoptera, Archostemata) in the Natural History Museum, London. *Paleontological Journal* 40 (1), 90–99.
- Ponomarenko, A.G., Ren, D., 2010. First record of *Notocupes* (Coleoptera: Cupedidae) in Locality Daohugou, Middle Jurassic of Inner Mongolia, China. *Annales Zoologici* 60 (2), 169–171.
- Pöthe De Baldis, E.D., 1984. Microfloras cenozoicas. In: Ramos, V.A. (Ed.), *Geología y Recursos Naturales de la Provincia de Rio Negro*. Relatorio del IX Congreso Geológico Argentino 2(4), 393–411.
- Quattrocchio, M., Del Papa, C., 2000. Paleoambiente de la Secuencia Maíz Gordo (Paleoceno tardio–Eoceno temprano?), Arroyo Las Tortugas, Cuenca del Grupo Salta (NO Argentina). *Palinología y Sedimentología. Revista Española de Paleontología* 15 (1), 57–70.
- Quattrocchio, M.E., Ruiz, L.C., 1999. Paleoambiente de la Formación Pedro Luro (Maastrichtiano?–Paleoceno) en base a palinomorfos, cuenca del Colorado, Argentina. *Ameghiniana* 36 (1), 37–47.
- Riccardi, A.C., 1984a. Las Asociaciones de Amonitas del jurásico y Cretácico de la Argentina. *Actas IX Congreso Geológico Argentino*, 559–595.
- Riccardi, A.C., 1984b. Las Zonas de Amonitas del Cretácico de la Patagonia (Argentina y Chile). *Memoria III Congreso Latinoamericano de Paleontología*, 394–405.
- Riccardi, A.C., 2002. Invertebrados del Cretácico Superior. En: *Geología y Recursos Naturales de Santa Cruz*. In: Haller, M.J. (Ed.), *Relatorio del XV Congreso Geológico Argentino*. El Calafate, 461–479.
- Riccardi, A.C., Rölleri, E.O., 1980. Cordillera Patagónica Austral. In: Leanza, A.F. (Ed.), *Geología regional Argentina*. Academia Nacional de Ciencias, Córdoba, 2, pp. 1173–1306.
- Riccardi, A.C., Aguirre Urreta, M.B., Medina, F., 1987. Aconeceratidae (Ammonitina) from the Hauterivian–Albian of southern Patagonia. *Palaeontographica* 196A, 105–185.
- Shcherbakov, D.E., 2007. An extraordinary new family of Cretaceous planthoppers (Homoptera: Fulgoroidea). *Russian Entomological Journal* 16, 139–154.
- Solórzano Kraemer, M.M., Petrulevičius, J.F., 2007. A new Plant Hopper (Insecta: Nogodinidae) from Chiapas amber, middle Miocene of Mexico. *Geobios* 40, 827–832.
- Soriano, C., Delclós, X., 2006. New cupedid beetles from the Lower Cretaceous of Spain and the palaeogeography of the family. *Acta Palaeontologica Polonica* 51, 185–200.
- Spicer, A.R., Rees, P.M., Chapman, J.L., 1994. Cretaceous phytogeography and climate signals. In: Allen, J.R.L., Hoskins, B.J., Sellwood, B.W., Spicer, R.A., Valdes, P.J. (Eds.), *Palaeoclimates and their Modelling; with Special Reference to the Mesozoic Era*. Chapman and Hall, pp. 69–77.
- Szweido, J., Azar, D., Nohra, Y., 2013. First record of Perforissidae from the Lower Cretaceous Lebanese amber (Hemiptera: Fulgoromorpha: Fulgoroidea). In: Azar, D., Engel, M.S., Jarzembowski, E., Krogmann, L., Nel, A., Santiago-Blay, J. (Eds.), *Insect Evolution in an Amberiferous and Stone Alphabet*. Brill, pp. 145–163.
- Tjallingii, W.F., 2006. Salivary secretions by aphids interacting with proteins of phloem wound responses. *Journal of Experimental Botany* 57, 739–745.
- Varela, A.N., 2011. Sedimentología y Modelos Deposicionales de la Formación Mata Amarilla, Cretácico de la Cuenca Austral, Argentina. Unpublished Ph.D. Thesis. Universidad Nacional de La Plata, Facultad de Ciencias Naturales y Museo, 287 pp.
- Varela, A.N., Richiano, S., Poiré, D.G., 2011. Tsunami vs storm origin for shell bed deposits in a lagoon environment: an example from the Upper Cretaceous of Southern Patagonia, Argentina. *Latin American Journal of Sedimentology and Basin Analysis* 18, 63–85.

- Varela, A.N., Poiré, D.G., Martin, T., Gerdes, A., Goin, F.J., Gelfo, J.N., Hoffmann, S., 2012a. U–Pb zircon constraints on the age of the Cretaceous Mata Amarilla Formation, Southern Patagonia, Argentina: its relationship with the evolution of the Austral Basin. *Andean Geology* 39 (3), 359–379.
- Varela, A.N., Veiga, D.G., Poiré, D.G., 2012b. Sequence stratigraphic analysis of Cenomanian greenhouse palaeosols: a case study from southern Patagonia, Argentina. *Sedimentary Geology* 271–272, 67–82.
- Varela, A.N., Gómez-Peral, L.E., Richiano, S., Poiré, D.G., 2013. Distinguishing similar volcanic source areas from an integrated provenance analysis: implications for foreland Andean basins. *Journal of Sedimentary Research* 83, 258–276.
- Wilckens, O., 1907. Die Lamellibranchiaten, gastropoden etc. der oberen Kreide Südpatagoniens. *Naturforschungen Gesellschaft Freiburg. Berichte* 15, 1–70.
- Zamuner, A.B., Poiré, D.G., Iglesias, A., Larriestra, F., Varela, A.N., 2004. Upper Cretaceous *in situ* Petrified Forest in Mata Amarilla Formation, Tres Lagos, Southern Patagonia, Argentina. Abstracts VII International Organization of Paleobotany Conference. Bariloche, p. 150.
- Zherikhin, V.V., 2002. 3.2. Ecological history of the terrestrial insects. In: Rasnitsyn, A.P., Quicke, D.L.J. (Eds.), 2002. *History of insects*. Kluwer Academic Publishers, pp. 331–387.