


Complete mitochondrial genome of the planthopper *Orthopagus splendens* (Germar, 1830) (Hemiptera: Fulgoromorpha: Dictyopharidae)

Yan-li Zheng^{a,b,c,d} , Thierry Bourgoïn^e , Lin Yang^{a,b} , Xiang-Sheng Chen^{a,b} , Xu-Qiang Luo^{c,d}  and Guang-Jie Luo^{c,d}

^aInstitute of Entomology, Guizhou University, Guiyang, PR China; ^bThe Provincial Key Laboratory for Agricultural Pest Management of Mountainous Region, Guiyang, PR China; ^cSchool of Geography and Resources, Guizhou Education University, Guiyang, PR China; ^dGuizhou Provincial Key Laboratory of Geographic State Monitoring of Watershed, PR China; ^eInstitut Systématique, Evolution, Biodiversité (ISYEB), MNHN-CNRS-Sorbonne Université-EPHE-Univ. Antilles, Museum National d'Histoire Naturelle, Paris, France

ABSTRACT

The first complete mitochondrial genome of a dictyopharid planthopper, *Orthopagus splendens* (Germar, 1830) (Hemiptera: Fulgoroidea: Dictyopharidae) is sequenced. The 15,349 bp long complete mitogenome contains 13 protein-coding genes (PCGs), 22 *tRNA* genes, two *rRNA* genes, and 1 A + T-rich region with an arrangement identical to that observed in most insect mitogenomes (GenBankNo. MW441850). All PCGs start with ATN, and end with TAN or single T (*nad1*, *nad5*, and *atp6*). A phylogenetic analysis places *O. splendens* as sister to Fulgoridae confirming a sister relationship between Dictyopharidae and Fulgoridae.

ARTICLE HISTORY

Received 15 June 2021
Accepted 29 July 2021

KEYWORDS

Orthopagus splendens;
Dictyopharidae; mitogenome; phylogeny

The planthoppers of the family Dictyopharidae Spinola, 1839 (Hemiptera: Fulgoromorpha), currently groups 740 species in 160 genera (Bourgoïn 2020). Divided into two subfamilies and 19 tribes, the species *Orthopagus splendens* (Germar, 1830) belongs to the Dictypharinae Spinola, 1839, Orthopagini Emeljanov 1983 (Muir 1923; Metcalf 1946; Song et al. 2018; Bourgoïn 2020). We report here the first complete mitochondrial genome (mitogenome) for this planthopper family.

Mitogenome DNA was extracted from a single female specimen collected from Xinliangtian Village, Guiding County, Guizhou province, China (107°23'E, 26°57'N) on the 11 August 2020. After DNA was extracted from the hind legs, a specimen was deposited at the Institute of Entomology, Guizhou University (Xiangsheng Chen, chenxs3218@163.com) under the voucher number GUGC-20200811-Y1. The mitogenome sequences were obtained by next-generation sequencing method (Illumina HiSeq X 10), assembled using Geneious version 10.2.2 (Kearse et al. 2012), annotated, and conducted using MITOZ (Meng et al. 2019) and deposited in GenBank, accession number MW441850. IQtree (Kumar et al. 2016) was used for phylogenetic analyses and to construct the maximum likelihood (ML) tree based on 13 PCGs, 22 *tRNA*, and two *rRNA* of 23 species (including four outgroups from

Cicadelloidea and Cicadoidea), and 18 related planthopper species from different Fulgoroidea families.

The complete mitogenome of *O. splendens* is 15,349 bp in length and consists of 13 protein-coding genes (PCGs), two ribosomal *rRNA* genes (*rRNAs*), 22 transfer *rRNA* genes (*tRNAs*), and one large non-coding region (D-loop: [A + T]-rich region). The D-loop part is 1181 bp, taking place between 12S *rRNA* and *trnI*. The overall bases composition is A: 30.2%, T: 47.2%, C: 14.6%, and G: 8%. AT skew $((A - T)/(A + T))$ and GC skew $((G - C)/(G + C))$ are 0.22, -0.294, respectively. All the 13 PCGs started with ATN, ended with TAN or a single T (*nad1*, *nad5*, and *atp6*) residue. The length of 22 *tRNA* ranged from 59 bp (*trnS*) to 70 bp (*trnK*). Genes of 16S *rRNA* and 12S *rRNA* are 1177 and 729 bp, respectively.

The phylogenetic tree obtained confirms the monophyly of Fulgoroidea and most nodes are recovered with the higher ultrafast bootstrap (UFB = 100). As expected from both morphological (Emeljanov 1979; Song et al. 2018) and molecular analyses (Urban and Cryan 2009; Song and Liang 2013), the dictyopharid species appears in a sister relationship with Fulgoridae, both in an unexpected clade sister to Achilidae: (Achilidae + (Dictyopharidae + Fulgoridae)). While already previously observed by Song and Liang (2013) and Wang et al. (2019), it is probably due to a limited sampling effect.




Figure 1. Maximum likelihood tree based on 23 complete mitochondrial genome sequences. Fulgoromorpha: 19 in-group species and four outgroup species: two in Cicadellidae and two in Cicadidae.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

This project was supported by the Program of Guizhou Science and Technology Foundation ([2019] 1242) and the National Natural Science Foundation of China [grant nos. 32060343 and 31472033].

ORCID

Yan-li Zheng <http://orcid.org/0000-0002-6964-5988>
 Thierry Bourgoin <http://orcid.org/0000-0001-9277-2478>
 Lin Yang <http://orcid.org/0000-0002-7841-5156>
 Xiang-Sheng Chen <http://orcid.org/0000-0001-9801-0343>
 Xu-Qiang Luo <http://orcid.org/0000-0003-3595-7446>

Data availability statement

The data supporting the findings of this study are openly available in the National Center for Biotechnology Information (NCBI) at <https://www.ncbi.nlm.nih.gov>, accession number MW441850. The associated BioProject, SRA, and Bio-Sample numbers are PRJNA726041, SRS8798664, and SAMN18920653, respectively.

References

Bourgoin T. 2020. FLOW (Fulgoromorpha Lists on the Web): a world knowledge base dedicated to Fulgoromorpha. Version 8. [accessed 4 May 2021]. <http://hemiptera-databases.org/flow/>.

- Emeljanov AF. 1979. Problema razgranicheniya semeïstv Fulgoridae i Dictyopharidae (Homoptera, Auchenorrhyncha). *Trudy Zool Inst Akademii Nauk SSSR*. 82:3–22.
- Kearse M, Moir R, Wilson A, Stones-Havas S, Cheung M, Sturrock S, Buxton S, Cooper A, Markowitz S, Duran C, et al. 2012. Geneious Basic: an integrated and extendable desktop software platform for the organization and analysis of sequence data. *Bioinformatics*. 28(12): 1647–1649.
- Kumar S, Stecher G, Tamura K. 2016. MEGA7: molecular evolutionary genetics analysis version 7.0 for bigger datasets. *Mol Biol Evol*. 33(7): 1870–1874.
- Meng G, Li Y, Yang C, Liu S. 2019. MitoZ: a toolkit for animal mitochondrial genome assembly, annotation and visualization. *Nucleic Acids Res*. 47(11):e63–e63.
- Metcalf ZP. 1946. General catalogue of the Hemiptera, Fasc. IV. Fulgoroidea, Part 8 Dictyopharidae. Northampton (MA): Smith College, Northampton, MASS; p. 246.
- Muir F. 1923. On the classification of the Fulgoroidea (Homoptera). *Proc Hawaiian Entomol Soc*. 205(5):247.
- Song N, Liang AP. 2013. A preliminary molecular phylogeny of planthoppers (Homoptera: Fulgoroidea) based on nuclear and mitochondrial DNA sequences. *PLoS One*. 8(3):e58400.
- Song ZS, Bartlett CR, O'Brien LB, Liang AP, Bourgoin T. 2018. Morphological phylogeny of Dictyopharidae (Homoptera: Fulgoromorpha). *Syst Entomol*. 43(4):637–658.
- Urban JM, Cryan JR. 2009. Entomologically famous, evolutionarily unexplored: the first phylogeny of the lanternfly family Fulgoridae (Insecta: Hemiptera: Fulgoroidea). *Molecular Phylogenetics and Evolution* 50: 471–484.
- Wang W, Huang Y, Bartlett CR, Zhou F, Meng R, Qin D. 2019. Characterization of the complete mitochondrial genomes of two species of the genus *Aphaena* Guérin-Méneville (Homoptera: Fulgoridae) and its phylogenetic implications. *Int J Biol Macromol*. 141:29–40.